

Magnetic field dependent transmission phase of a double dot system in a quantum ring

M. Sigrist,¹ A. Fuhrer,¹ T. Ihn,¹ K. Ensslin,¹ S. E. Ulloa,^{1,2} W. Wegscheider,³ M. Bichler⁴

¹*Solid State Physics, ETH Zürich, 8093 Zürich, Switzerland*

²*Department of Physics and Astronomy, Ohio University, Athens, Ohio 45701-2979*

³*Institut für experimentelle und angewandte Physik, Universität Regensburg, Germany*

⁴*Walter Schottky Institut, Technische Universität München, Germany*

(Dated: February 2, 2008)

The Aharonov-Bohm effect is measured in a four-terminal open ring geometry based on a Ga[Al]As heterostructure. Two quantum dots are embedded in the structure, one in each of the two interfering paths. The number of electrons in the two dots can be controlled independently. The transmission phase is measured as electrons are added to or taken away from the individual quantum dots. Although the measured phase shifts are in qualitative agreement with theoretical predictions, the phase evolution exhibits unexpected dependence on the magnetic field. For example, phase lapses are found only in certain ranges of magnetic field.

In mesoscopic systems the conductance and the quantum mechanical transmission of charge carriers between contacts are intimately related. The corresponding transmission amplitude is a complex number with magnitude and phase. The phase difference of interfering paths in coherent quantum rings can be detected in the conductance by measuring Aharonov-Bohm (AB) oscillations. Keeping the transmission phase of one of the interfering paths constant (reference path), the phase change of the other can be measured. This technique opens the attractive possibility to investigate the transmission phase which contains information about the system complementary to the transmission probability. Multiterminal devices are required for such experiments in order to circumvent the generalized Onsager symmetry relations [1] restricting the transmission phase to values of 0 or π in two-terminal devices.

Ring and ring-like interference geometries have been exploited for many types of experiments [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12]. The electronic phase in particular was studied in a series of pioneering publications [2, 3, 4, 5, 6]. The transmission phase through a two-terminal device with a single quantum dot has been investigated [2] and it was concluded that electron transport through a Coulomb blockaded quantum dot is at least partially phase coherent. The phase of the reflection coefficient of a quantum dot in the integer quantum Hall regime was measured in Ref. 3. Later, the transmission phase of a quantum dot was studied in a multi-terminal configuration [4]. It was found that the transmission phase changes by about π across a typical Coulomb blockade resonance, as expected. Between resonances the experiment exhibits so-called phase lapses rather than the almost constant phase expected in the simplest model. Similar experiments were carried out on a Kondo-correlated system in Refs. 5 and 6.

A large number of theoretical papers (for a review see [13]) has addressed the issue of the phase lapses and suggested possible explanations. It was argued that the transmission phase has to be distinguished from the Friedel phase which fulfills a general sum rule and is a monotonous function of energy, while the transmission

phase can show phase lapses of π when the transmission goes through zero [14, 15, 16]. It was also pointed out that the measured phase may depend on the details of the entire interferometer, on interactions [17], and on the dot-lead coupling [18]. The phase determined experimentally is the transmission phase through the dot, only if the system is unitary [19], i.e. it couples negligibly to its environment. A tunneling-Hamiltonian approach predicted the phase to be influenced by the number and width of leads connecting to the ring [20, 21]. In a scattering matrix approach, however, the limit of many occupied modes in the channels reflects the energy dependence of the resonance phase shift [22]. The equivalence of the tunneling-Hamiltonian formulation and the scattering matrix approach are discussed in [23]. Theoretical proposals have considered the situation of two quantum dots in an AB interferometer and the possible coupling of the dots via phase coherent transport through the leads [24, 25]. In the context of spintronics two coupled quantum dots in an AB interferometer may be used for detecting entanglement of spin states [26].

Here we investigate the phase evolution of a system of two quantum dots with negligible mutual electrostatic interaction embedded in two arms of a four-terminal AB ring. Phase measurements in this unique system are desirable, because both arms of the ring including the two dots can be tuned individually, but the phase coherence of the entire system provides an inherent coupling mechanism [24] and makes transport through the dots interdependent. The arrangement enables us to control and analyze the effect of the interferometer reference arm on the measured phase experimentally. The phase evolution of the system is studied when a single electron is added to either of the two quantum dots, to both quantum dots simultaneously, or when an electron is transferred from one dot to the other. The observed phase shifts agree qualitatively with theoretical expectations at elevated magnetic fields. Unexpectedly, the phase evolution is found to depend on the magnetic field and occasional phase lapses occur in certain field ranges. These observations highlight the need to consider nonlocal coherent effects in the entire system.

The sample is a Ga[Al]As heterostructure containing a two-dimensional electron gas 37 nm below the surface. The lateral pattern was fabricated with the biased tip of an atomic force microscope which locally oxidizes the GaAs surface. Details of this fabrication technique are described in Ref. 27. Figure 1(a) shows a micrograph of the oxidized pattern. Lateral gate electrodes marked pg1 through pg4 are used to tune the conductance in each of the four quadrants of the ring. All measurements were carried out at 100 mK in a dilution refrigerator.

The ring was characterized in the open regime where each quadrant supports 2-4 lateral modes. A pronounced AB effect is observed in a local measurement setup in which the current and voltage contacts are the same two terminals of the ring [28]. The AB signal is maximized with a non-local setup: a bias voltage V_{bias} was applied to terminal A [see Fig. 1(b)]. The lower and upper contacts (B and D) were grounded via current-voltage converters measuring the currents I_B and I_D . The non-local voltage V_{nl} was measured at terminal C. Figure 1(c) shows the AB oscillations in V_{nl} . Their period $\Delta B = \Phi_0/A = 4.8$ mT is consistent with the area $A = 0.85 \mu\text{m}^2$ of the ring.

If any individual segment is tuned close to pinch off with a negative voltage applied to the corresponding plunger gate, a quantum dot is induced exhibiting Coulomb blockade. As an example, Fig. 1(d) shows the




FIG. 1: (a) AFM micrograph of the ring structure. Oxide lines (bright lines) fabricated by AFM lithography lead to insulating barriers in the two-dimensional electron gas. Areas marked pg1-pg4 are used as lateral gates to tune the conductance of the four arms of the ring and eventually act as plunger gates for the dots. (b) Schematic arrangement of the ring with its four terminals (c) Non-local voltage detected in the open regime of the ring. (d) Conductance through a quantum dot induced in segment 1 as a function of plunger gate voltage. Here, segment 3 was completely pinched off.

conductance of segment 1 [A to D in Fig. 1(b)] as a function of V_{pg1} with segment 3 completely pinched off and segments 2 and 4 open. Clear Coulomb blockade oscillations are observed. The location of the quantum dot can be estimated from the lever arms of gates pg1 to pg4. The dots form within the segments and not at the openings to the contacts. From Coulomb blockade diamonds measured as a function of forward bias we find a typical Coulomb charging energy of 1 meV, which corresponds in a disc capacitor model roughly to the area of a single segment. More details about the characterisation of the ring can be found in Ref. 28.

We proceed to study Coulomb blockade in the measurement configuration of Fig. 1(a). For the following experiments we have carefully chosen a series of conductance peaks of one well defined quantum dot in segment 1 (dot 1) and another one in segment 2 (dot 2). Segments 3 and 4 are open. The current signals I_B and I_D are shown in color as a function of the gate voltages V_{pg1} and V_{pg2} in Figs. 2b and c. The current I_B (I_D) shows predominantly the Coulomb maxima of the dot in segment 2 (segment 1) while those of the other dot are much weaker. The non-local voltage in Fig. 2(a) displays clear signatures of conductance resonances of both quantum dots. The lever arm of gate voltage 1 (2) on the dot in segment 2 (1) is about a factor of six smaller than for the direct gate voltage. There are stripes with an intermediate slope in all three quantities which we attribute to the formation of standing waves between the two quantum dots near lead A. This resonator is coupled in series to




FIG. 2: Color plots of (a) the non-local voltage V_{nl} , (b) the current I_B and (c) I_D as a function of the two plunger gate voltages V_{pg1} and V_{pg2} which tune the quantum dots in segments 1 and 2 of the ring.




FIG. 3: AB oscillations in non-local voltage V_{nl} as function of the plunger gate voltages tuned along trace ‘linear2’. (a) The currents I_D and I_B at a magnetic field of 95 mT. (b) Color plot of V_{nl} as function of plunger gates and magnetic field. (c) Cross-sections along dashed lines in (b) giving V_{nl} vs. magnetic field. The dashed lines mark constant values of magnetic field where a π phase shift of the AB oscillations can be seen, as gate voltages are tuned across the conductance peak in dot 2.

both quantum dots, a fact which leads to a corresponding modulation of the dot currents in certain ranges of gate voltages and magnetic fields.

In order to study the phase evolution of the transmission through the quantum dots we measure V_{nl} as a function of magnetic field. The results presented below were obtained in a regime where the coupling to the leads was strong enough, but the dots were still well defined. By tuning V_{pg1} and V_{pg2} we follow the trace labeled ‘linear2’ in Fig. 2(a), staying on a conductance maximum in dot 1 while stepping through a conductance peak in dot 2. As shown in Fig. 3(a) the current I_B shows the expected Coulomb peak of dot 2 as a function of plunger gate voltage, while I_D changes little. At each gate voltage setting the AB oscillations are measured. The non-local voltage in Fig. 3(b) shows a pronounced AB signal which is strongest in the range of the conductance maximum. A set of specific traces of V_{nl} versus magnetic field is depicted in Fig. 3(c). The vertical lines connect minima in the lowest trace with maxima in the uppermost trace. This corresponds to a phase shift of π .

The multi-probe conductance formula [1] gives for the non-local voltage in our setup:

$$V_{nl} = -\frac{T_{CA}}{T_{CC}} V_{bias}.$$

The transmission coefficients T_{ij} from probe j to i are named according to Fig. 1(b). Considering only oscillations with a periodicity of h/e we assume $T_{CC} = -\sum_{i \neq C} T_{Ci}$ to be constant. In this case AB oscillations in V_{nl} reflect the interference contribution to T_{CA} from which the phase difference between the two interfering transmission amplitudes can be directly read [4]. In contrast, phase changes seen in the two currents do not di-




FIG. 4: AB oscillations along the different traces indicated in Fig. 2. (a) sweep ‘linear 2’ (b) sweep ‘diag 1’ (c) sweep ‘linear 1’ and (d) sweep ‘diag 2’. Vertical lines are guides to the eye. Dashed horizontal lines indicate, where maxima in the magnetic field averaged contribution to V_{nl} occur.

rectly reflect the difference of transmission phases of the dots. In the following we therefore consider only phase shifts of the AB oscillations in V_{nl} .

We have analyzed such phase shifts quantitatively for all four traces indicated by the arrows in Fig. 2(a). The fast Fourier transform of each magnetic field sweep was multiplied with the filter function $f(\omega) = (\sigma\omega)^2/2 \exp[1 - (\sigma\omega)^2/2]$ with $\sigma = h/(e\sqrt{2}\pi A)$ in order to obtain the pure h/e -periodic contribution. The inverse fast Fourier transform of the filtered data gives the oscillatory component of V_{nl} as a function of magnetic field which is plotted in Fig. 4. We have verified that this filtering procedure does not influence the phase of the AB oscillations by carefully comparing filtered data with raw data.

We begin the discussion with Fig. 4(a) corresponding to trace ‘linear2’ in Fig. 2(a). The dashed horizontal line indicates the maximum in V_{nl} averaged over B . The AB amplitude is strongest close to this line, as expected, but displays a pronounced B -dependence, i.e. it is significantly larger for $B > 30$ mT than below. The phase shift can be directly read from the shift of AB maxima or minima as a function of V_{pg2} . The general trend is a shift to larger magnetic fields with increasing V_{pg2} . The expected phase shift of π across the conductance resonance is clearly discernible for $B > 50$ mT. However, there is a dip in the AB amplitude at $V_{pg2} \approx -68$ mV which has a strong phase shift associated with it. At $B \approx 40$ mT and

$V_{pg2} \approx -68$ mV [arrow in Fig. 4(a)] the AB amplitude is zero and a phase lapse of π can be identified. This phase lapse lies in the flank of the peak of the field averaged V_{nl} and occurs at a point of zero AB amplitude, similar to the phase lapses observed in Ref. 4 and in agreement with the general theoretical discussion of phase lapses in Ref. 15. The novel aspect here is the occurrence of the phase lapse only in a limited range of magnetic field. Beyond this field range the phase lapse disappears in favor of a continuous phase evolution of order π .

Along trace ‘diag1’ in Fig. 2(a) an electron is added to both dots simultaneously. In this case we expect a phase shift of $\pi - \pi = 0$ since the phase difference between the arms of the ring remains constant. In the experiment the phase shift in Fig. 4(b) is indeed small for $B > 50$ mT, it amounts to about $\pi/10$. At lower fields, however, phase lapses and related strong phase shifts occur like that observed for trace ‘linear2’.

When we measure along trace ‘linear1’ [Fig. 4(c)] the gross trend of the phase shift differs in sign from that in Fig. 4(a), in agreement with the fact that we tune the dot in the other segment. In the coupling regime of the quantum dots where AB oscillations are observable, Coulomb resonances can occur close to each other. This is the case for this sweep as indicated by the two dashed blue horizontal lines in Fig. 4(c). In such a case the question arises how the transmission phases of individual resonances combine to the observed phase shift. In our measurement we find that the phase accumulation across the first resonance just above $V_{pg1} = -60$ mV is about $-\pi$ for B between 30 and 70 mT, as expected. At higher fields the phase shift weakens. At $V_{pg1} \approx -55$ mV and between -40 and -50 mV, phase lapses occur which influence the phase evolution at higher magnetic fields but cannot compensate the accumulated phase completely. At the top edge of the figure, the second resonance leads to a further phase accumulation in the same direction as the first.

Along trace ‘diag2’ [Fig. 4(d)] an electron is removed from dot 2 and added to dot 1. As expected, AB oscillation maxima shift in the same direction as in ‘linear1’. However, at $B = 80$ mT, for example, the phase accumulated between $V_{pg1} = -68$ mV and -56 mV is only slightly more than π rather than 2π . Again, the phase evolution is influenced by occasional phase lapses.

Summarizing the observations we can make the following statements. We have succeeded in measuring the AB phase in a four terminal quantum ring with two dots embedded in two different segments. The AB oscillations tend to be well developed and suitable for phase measurements only at slightly elevated magnetic fields. In the range between 50 and 80 mT the expected phase shifts across Coulomb-blockade resonances are typically observed. Phase lapses occur occasionally in finite magnetic field ranges at specific gate voltages. They are accompanied by a vanishing AB amplitude. Even outside the B -ranges where phase lapses occur, the phase evolution can be strongly modified by their presence. Fol-

lowing the argument of Büttiker and coworkers [15, 16], the transmission amplitude follows a certain curve in the complex plane as a function of gate voltage. A phase lapse accompanied with a zero in transmission occurs when this line goes through the origin. A change in magnetic field may shift this curve a little leading to a strong, but continuous phase shift accompanied with small transmission. The appearance of phase lapses in finite magnetic field ranges impedes the definition of a magnetic field independent transmission phase. A conceivable origin of the phase lapses is the finite width of the ring segments accommodating several modes. In the magnetic field range investigated, the classical cyclotron radius is larger or comparable to the ring radius. A small influence of Lorentz force effects cannot be excluded at the highest fields.

Our samples differ from those in Ref. 4 by design and in view of the technological approach. We have adopted the general idea of using reflecting walls to guide the electrons ballistically around the ring without losing too many of them into the contacts. Compared to Ref. [4] the most noticeable feature of our structure is the second dot in the reference arm. Its addition gives individual control over the accumulated phases in both arms. Since the reference dot is kept on a conductance resonance, i.e. in the position of maximum slope of the phase evolution, while the other is swept through a resonance, the reference dot is very sensitive to phase changes. Should the reference dot be tuned slightly off its resonance during the sweep ‘linear 2’, one would expect a pronounced influence on the measured phase evolution. The fact that we still observe a phase change of about π on trace ‘linear2’, when dot 2 is tuned through a conductance resonance, lets us conclude that the transmission phase through dot 1 is rather stable and the observed phase shift is dominated by dot 2.

In this experiment, the transmission phase was measured in a system where two Coulomb blockaded quantum dots are embedded in a four-terminal quantum ring. We have shown that the phase evolution can be determined with individual control over the electron occupancy in each dot. The experiments demonstrate that the measured transmission phase depends on the magnetic field range analyzed. Phase lapses can arise at certain magnetic fields, while at others they give way to a continuous phase evolution. In certain magnetic field ranges the phase behavior is in agreement with theoretical expectations, but its behavior is much more complex than anticipated. This possibly suggests that structural resonances, in addition to the dots, play a role in the overall transmission phase of the system.

Acknowledgments

We thank M. Büttiker, Y. Gefen, M. Heiblum, H. Weidenmüller for valuable discussions. Financial support by the Swiss Science Foundation (Schweizerischer National-

fonds) is gratefully acknowledged.

-
- [1] M. Büttiker, Phys. Rev. Lett. **57**, 1761 (1986).
 - [2] A. Yacoby, M. Heiblum, D. Mahalu, and H. Shtrikman, Phys. Rev. Lett. **74**, 4047 (1995).
 - [3] E. Buks, R. Schuster, M. Heiblum, D. Mahalu, V. Umansky, and H. Shtrikman, Phys. Rev. Lett. **77**, 4664 (1996).
 - [4] R. Schuster, E. Buks, M. Heiblum, D. Mahalu, V. Umansky, and H. Shtrikman, NATURE **385**, 417 (1997).
 - [5] Y. Ji, M. Heiblum, D. Sprinzak, D. Mahalu, and H. Shtrikman, SCIENCE **290**, 779 (2000).
 - [6] Y. Ji, M. Heiblum, and H. Shtrikman, Phys. Rev. Lett. **88**, 076601 (2002).
 - [7] E. Buks, R. Schuster, M. Heiblum, D. Mahalu, and V. Umansky, NATURE **391**, 871 (1998).
 - [8] D. Sprinzak, E. Buks, M. Heiblum, and H. Shtrikman, Phys. Rev. Lett. **84**, 5820 (2000).
 - [9] K. Kobayashi, H. Aikawa, S. Katsumoto, and Y. Iye Phys. Rev. Lett. **88**, 256806 (2002).
 - [10] A. W. Holleitner, C. R. Decker, H. Qin, K. Eberl, and R. H. Blick Phys. Rev. Lett. **87**, 256802 (2001).
 - [11] A. W. Holleitner, R. H. Blick, A. K. Hüttel, K. Eberl, and J. P. Kotthaus, SCIENCE **297**, 70 (2002).
 - [12] A. Fuhrer, S. Lüscher, T. Ihn, T. Heinzel, K. Ensslin, W. Wegscheider, and M. Bichler, Nature **413**, 822 (2001); U.F. Keyser, C. Fühner, S. Borck, R.J. Haug, M. Bichler, G. Abstreiter, W. Wegscheider, Phys. Rev. Lett. **90**, 196601 (2003).
 - [13] G. Hackenbroich, Phys. Rep. **343**, 463 (2001).
 - [14] H.-W. Lee, Phys. Rev. Lett. **82**, 2358 (1999).
 - [15] T. Taniguchi and M. Büttiker, Phys. Rev. B **60**, 13814 (1999).
 - [16] A.L. Yeyati and M. Büttiker, Phys. Rev. B **62**, 7307 (2000).
 - [17] J. König and Y. Gefen, Phys. Rev. B **65**, 045316 (2002).
 - [18] A. Silva, Y. Oreg, and Y. Gefen, Phys. Rev. B **66**, 195316 (2002).
 - [19] O. Entin-Wohlman, A. Aharony, Y. Imry, Y. Levinson, and A. Schiller, Phys. Rev. Lett. **88**, 166801 (2002).
 - [20] A. Aharony, O. Entin-Wohlman, B. I. Halperin, and Y. Imry, Phys. Rev. B **66**, 115311 (2002).
 - [21] Y. Gefen, private communication
 - [22] H. A. Weidenmüller, Phys. Rev. B **65**, 245322 (2002).
 - [23] B. Kubala and J. König, cond-mat /0212536
 - [24] J. König and Y. Gefen, Phys. Rev. B **65**, 045316 (2002)
 - [25] B. Kubala and J. König, Phys. Rev. B **65**, 245301 (2002)
 - [26] D. Loss and E. V. Sukhorukov, Phys. Rev. Lett. **84**, 1035 (2000).
 - [27] R. Held, S. Lüscher, T. Heinzel, K. Ensslin, and W. Wegscheider, Appl. Phys. Lett. **75**, 1134 (1999), for a review see A. Fuhrer, A. Dorn, S. Lüscher, T. Heinzel, K. Ensslin, W. Wegscheider, and M. Bichler, Superl. Microstr. **31**, 19 (2002).
 - [28] M. Sigrist, A. Fuhrer, T. Ihn, K. Ensslin, W. Wegscheider, M. Bichler, cond-mat/0307269.